

Mosquito Surveillance Report

Joe Brew and Ben Brew

October 14, 2014

Contents

Executive Summary	2
Visual Overview	3
Time	3
Space	3
Normality	3
Disease Vectors	3
Forecast	4
Vectors of Disease by Location	5
Spatial interpolation maps	6
Mosquito Types	7
Details of Predictive Model	9



Executive Summary

Most Recent Collection

As predicted, mosquito numbers remained low over recent weeks, ranging from 105 to 131 per trap through the most recent collection (October 02, 2014).

Forecast

We forecast that mosquito levels will remain moderate/low through the middle of next week (90% confidence interval of 0-352 through October 22).


Predictive Model Validation

At 131 specimens per trap, the most recent collection was similar to our prediction of 123, and well within the 90% confidence interval for the range of prediction.

Visual Overview

Time

The current mosquito population is lower than usual for this time of year (and significantly lower than last year's mid-September spike).




Space

Mosquitoes were largely scattered throughout the county, with a slight concentration in the northern traps.




Normality

The most recent collection was at levels equivalent to approximately the 50th percentile of historical (2008-13) levels.







Disease Vectors

Mosquitoes capable of carrying WNV are much more predominant now than at the beginning of the summer.



Forecast

We use recursive, quadratic linear regression modelling to forecast the average number of mosquitoes per trap up to 15 days in advance.¹



¹We are actively experimenting with non-parametric approaches to improve modelling accuracy, and expect to have a modified KNN model with better results by late September.


Vectors of Disease by Location

Mosquito species capable of carrying WNV and SLEV were most prevalent in recent trap collections.




Spatial interpolation maps


Mosquito species cabale of carrying WNV and SLEV were most prevalent in recent trap collections.



Mosquito Types

No species saw abnormal increases in recent weeks.





Details of Predictive Model

Historically, the model has performed well, correctly predicting the late summer spikes in 2012 and 2013. Given the preference for accuracy at high numbers, the model intentionally includes outlying high observations, thereby weighting them.

Having simulated more than 65,000 unique models, our best fit equation (using the sum of least squares approach) was:

$$\hat{Y} = \beta_0 + \beta_1^2(5.6508) + \beta_2(0.5938)$$

where \hat{Y} is the estimated mean number of mosquitoes per trap, β_0 is set to 0, β_1 is the cumulative rainfall in the period 15 to 29 days prior to the date of prediction and β_2 is the mean number of mosquitoes per trap in the most recent prior trap collection.

Though an original model relied only on rainfall, incorporating the most recent trap prediction saw our R-squared improve from 0.52 to 0.82. This means that we can now explain over 80% of the variance in mosquito populations up to 15 days ahead of time.